

Computational Accelerator Physics at NERSC: A User's Perspective

Robert Ryne

Lawrence Berkeley National Laboratory


Outline

- Overview of accelerators and the computational accelerator physics community
- The types of calculations we do at NERSC
- Personal observations about NERSC
- Future challenges in accelerator S&T and the need for HPC

The National Investment in Particle Accelerators is Enormous



Lawrence Berkeley Natl. Lab.
ALS



Indiana Univ.
IUCF



Fermi National Acc. Lab.
Tevatron



Michigan State Univ.
NSCL



Argonne National Lab.
IPNS ATLAS APS



Stanford Linear Acc. Center
SLC PEP-II SSRL



Cornell Univ. **CESR CHSS**



Los Alamos Natl. Lab.
LANSCE/Lujan



Oak Ridge National Lab.
SNS






Thomas Jefferson National
Accelerator Facility: **CEBAF**





Brookhaven National Lab.
RHIC NSLS

Contributions of accelerators have significant economic impact and greatly benefit society

- Medical isotope production
- Electron microscopy
- Accelerator mass spectrometry
- Medical irradiation therapy
- Ion implantation
- Beam lithography




- Transmutation of waste
- Accelerator-driven energy production
- Hydrodynamic imaging




The accelerator community uses high performance computing to tackle of wide range of problems


- Designing complicated electromagnetic structures



- Modeling high intensity beam dynamics




- Exploring beams under extreme conditions



Why do we need terascale computing for next-generation accelerator design?

- High accuracy requirements
 - Design of 3D electromagnetic components
 - frequency accuracy to 1:10000
- Large-scale requirements
 - Designing 3D electromagnetic components
 - system-scale modeling
 - Modeling 3D intense beam dynamics
 - Halos, beam-beam effects, circular machines
 - Modeling 3D advanced accelerator concepts
 - laser- and plasma-based accelerators
- More physics
 - collisions, multi-species, surface effects, ionization, CSR, wakes,...

Accelerating Scientific Discovery in Accelerator Technology and Beam Physics: A SciDAC Multi-disciplinary, Multi-institutional Collaboration




Noteworthy computations that we have performed at NERSC

- First successful 3D eigenmode calculation of ~50 cell Next Linear Collider (NLC) structure including accelerating cells and damping manifold
- First self-consistent 3D Fokker-Planck simulation
- Simulations identified possible heating mechanism in PEP-II B-factory interaction region
- Simulations in support of SLAC E-157 plasma wakefield accelerator experiment
- Simulations in support of Spallation Neutron Source linac design effort
- Simulations in support of CERN/SPL design effort (similar to US neutrino factory design effort)

Challenges in Electromagnetic Systems Simulation: Example – NLC Accelerator Structure (RDDS) Design




- Start w/ cylindrical cell geometry
- adjust geometry for maximum efficiency
- add micron-scale variations from cell-to-cell to reduce wakefields
- stack into multi-cell structure
- Add damping manifold to suppress long-range wakefields, improve vacuum conductance, but preserve RDS performance. *Highly 3D structure.*



Require 0.01% accuracy
in accelerating frequency to
maintain structure efficiency
(**High resolution modeling**)




Verify wake suppression
in entire 206-cell section
(**System scale simulation**)




Parallel solvers needed to model large, complex 3D electromagnetic structures to high accuracy

NLC RDDS 47 Cell Stack - *Omega3P*

Calculation of modes in entire structure has begun




1st convergence of a mode in a dense spectrum



3D First-Principles Fokker-Planck Simulation

- Requires analog of 1000s of space-charge calculations/step
 - “...it would be completely impractical (in terms of # of particles, computation time, and statistical fluctuations) to actually compute [the Rosenbluth potentials] as multiple integrals” J.Math.Phys. 138 (1997).
- Feasibility demonstrated!

Self-Consistent
Diffusion
Coefficients




← Spitzer
approximation


Previous approximate
calculations performed
w/out parallel computation
were not self-consistent

PEP II - IR Beamline Complex


Left crotch




Center beam pipe




Right crotch



Identify localized modes to understand beam heating




Short section from IP



Laser/Plasma-based acceleration can produce gradients ~ 100 GeV/m



- High gradients measured in the lab over short distances
 - 100s to 1000s times greater than conventional technology
- **Plasma wakefield accelerator** (PWFA) concept uses extremely high fields in plasmas





- **Challenge is to control & stage** high-gradient sections to produce a high quality, high energy beam
- Simulation of 1-10 GeV PWFA would require $\sim 10\text{K}-100\text{K}$ CPU-hours

Direct comparison between expt and full-scale PIC simulation using OSIRIS
of the refraction of a 30 GeV e^- beam at a plasma vacuum interface

Experiment




PIC Simulation




Linac Modeling: Old vs. New Capability

- 1980s: **10K** particle, 2D serial simulations
- Early 1990s: **10K-100K**, 2D serial simulations
- 2000: **100M** (~10 hrs on 256 PEs); *more realistic model*





LEDA halo expt; 100M particles




SNS linac; 500M particles

Future Challenges in Beam Dynamics Modeling

- **Intense beams in circular machines** is a major challenge:
 - 100 to 1000 times more challenging than linacs
- Additional physics adds further complexity
 - Intense beams in bending magnets
 - Collisions
 - Wakefields
 - Radiation
 - Ionization



“banana” effect in circular machine code



Personal Observations

Personal observations about NERSC

- Outstanding support by User Services
 - Reliable, timely, friendly
 - Great phone support
 - Excellent web pages
 - Ease of adding new users
- Science-enabling resources
 - Seaborg, HPSS
 - ESnet
 - Auxiliary systems (viz, math servers)
 - Stable environment
- Algorithm and software development
 - Linear solvers, eigensolvers, cluster development,...
- Outstanding support staff
 - Event coordinators, training, publications,...

Personal observations, cont.

- NERSC does an excellent job of serving the needs of **both** a **large user community** and a **small number of power users/projects**
- NERSC has made the right choices in hardware. The staff has been very successful at procuring systems that are high-end yet with the stability and reliability to serve its large community of users
- NERSC provides enhanced support to selected projects
 - P.O.C. for the SciDAC accelerator modeling project
 - Server support, cvs, software requests, technical support,...

Future Activities in Accelerator S&T and the Need for HPC


Particle accelerators are among the most broadly applicable, most important scientific instruments in the world

We may be on the brink of major new discoveries



Given the complexity and importance of accelerators, it is imperative that the most advanced HPC tools be brought to bear on their design, optimization, commissioning, and operation

Accelerators are Crucial to Scientific Discoveries in High Energy Physics, Nuclear Physics, Materials Science, and Biological Science


“Starting this fall, a machine called RHIC will collide gold nuclei with such force that they will melt into their primordial building blocks”



“A new generation of accelerators capable of generating beams of exotic radioactive nuclei aims to simulate the element-building process in stars and shed light on nuclear structure”



“Biologists and other researchers are lining up at synchrotrons to probe materials and molecules with hard x-rays”



“Violated particles reveal quirks of antimatter”

“Muon Experiment Challenges Reigning Model of Particles”



Particle accelerators are among the largest,
most complex, most expensive scientific
instruments in the world

Cost of leading-edge high energy accelerators: several billion \$
Design decisions may have huge consequences

Example: Impact of Simulation



- Without large-scale simulation: **cost escalation**
 - Superconducting Super Collider (SSC): 1 cm increase in aperture due to lack of confidence in design resulted in \$1B cost increase
- With large-scale simulation: **cost savings**
 - \$100M reduction is estimated cost due to improved accelerator structure design for the Next Linear Collider (NLC)

Ongoing/Planned Simulation Activities


- PEP-II (heating)
- FNAL booster, BNL AGS, LANL PSR (beam loss)
- PEP-II, Tevatron, RHIC (beam-beam effect, luminosity)
- Next Linear Collider
- Neutrino factory/muon collider
- SLAC E162 (plasma wakefield accelerator)
- Laser driven accelerators (I'OASIS, UMichigan, UCLA, NRL, UTexas, LLNL,...)
- Next-generation light source

Opportunities at Next-Generation Accelerator Facilities

Exploring physics beyond the Standard Model. Are there new particles? New interactions?




Research with exotic nuclei: The nature of nucleonic matter; origin of the elements; tests of the Standard Model



Experimental Areas:
1: < 12 MeV/u 2: < 1.5 MeV/u 3: Nonaccelerated 4: In-flight fragments

Research using intense, ultra-short pulses of x-ray radiation (4th generation light source): fundamental quantum mechanics; atomic, molecular, and optical physics; chemistry; materials science; biology



HPC will play a major role

- **Present accelerators:** Maximize investment by
 - optimizing performance
 - expanding operational envelopes
 - increasing reliability and availability
- **Next-generation accelerators**
 - better designs
 - feasibility studies
 - Facilitate important design decisions
 - completion on schedule and within budget
- **Accelerator science and technology**
 - help develop new methods of acceleration
 - explore beams under extreme conditions

